References

Abdul Khaliq, Amar Matloob, Zahid Ata Cheema, and Muhammad Farooq,(2011). Allelopathic activity of crop residue incorporation alone or mixed against rice and its associated grass weed jungle rice (*Echinochloa colona* [l.] link), Chilean Journal of Agri. Res., 71(3).

Abdul Khaliq, Muhammad Yasir Riaz and Amar Matloob, (2013). Bio-economic assessment of chemical and nonchemical weed management strategies in dry seeded fine rice (*Oryza sativa* L.), African Journal of Agronomy, 1(1), 010-016.

Ademiluyi Benson Oluwafemi., (2014). Allelopathic effects of *Moringa olifera* on the germination and seedling survival of *Euphorbia heterophylla* L., G.J.B.A.H.S., 3(1) : 195-198.

Callaway, R.M., Nadkarni, N.M. and Mahall, B.E., 2002. Facilitation and interference of *Quercus douglasii* on understory productivity in central California. Ecology 72, 1484-1499.

Chou, C.H., 1999. Methodologies for allelopathy research: from fields to laboratory. Recent Advances in allelopathy. vol.I. A science for the future 1, 3-24.

De Candolle, A.P. (1832). Physiologie Vegetale. Bechet Jeune, Paris, France.

El-Rokiek, Kowthar G; Rafat. R. El-Masry; Nadia K. Messiha and Salah A. Ahmed, (2010). The Allelopathic Effect of Mango Leaves on the Growth and Propagative Capacity of Purple Nutsedge (*Cyperus rotundus* L.) Journal of American Science : 6(9).

Filemon Elisante and Patrick A. Ndakidemi (2014). Allelopathic effect of *Datura stramonium* on the survival of grass and legume species in the conservation areas , American Journal of Research Communication www.usa-journals.com Elisante, *et al*., 2014: Vol 2(1) 27.

Hamid Nouri, Zabiholah Ansari Talab, Abolfazl Tavassoli, (2012). Effect of weed allelopathic of sorghum (*Sorghum halepense*) on germination and seedling growth of wheat, Alvand cultivar. Annals of Biological Research, 3 (3):1283-1293.

Hamid Reza Miri (2011). Allelopathic Potential of Various Plant Species on Hordeum Spontaneum. Advances in Environmental Biology, 5(11): 3543-3549.

Hussain, S., Siddiqui, S.U., Khalid, S., Jamal, A., Qayyum, A. and Ahmad, Z., 2007. Allelopathic potential of Senna (*Cassia angustifolia* Vahl.) on germination and seedling characters of some major cereal crops and their associated grassy weeds. Pakistan Journal of Botany 39(4), 1145-1153

Indrejit, I., 2001. Environnemental effects on allelochemical activity. Agronomy Journal 93(1), 79-84.

J. Kayode and J. M. Ayeni, (2009). Allelopathic Effects of Some Crop Residues on the Germination and Growth of Maize (Zea mays L),The Pacific Journal of Science and Technology, 10(1).

J.A. Dhole, K.D. Lone , G.A. Dhole and S.S. Bodke (2013).Allelopathic effect of aqueous and ethanolic extracts of some common weeds on seed health of *Triticum aestivum* L. (Wheat), Int. J. Curr. Microbiol. App. Sci., 2(6): 254-260.

Jabran, K., Cheema, Z.A., Farooq, M. and Khaliq, A., 2007. Evaluation of fertigation and foliar application of some fertilizers alone and in combination with allelopathic water extracts in wheat. Proc International Workshop on Allelopathy – Current Trends and Future Applications 24, 18–21

Jai Knox, Disha Jaggi & M.S. Paul (2010). Evaluation of allelopathic potential of selected plant species on *Parthenium hysterophorus*, Egyptian Journal of Biology, 12, 57-64.

Javaid Iqbal, Zahid Ata Cheema and M. Naeem Mushtaq (2009). Allelopathic Crop Water Extracts Reduce the Herbicide Dose for Weed Control in Cotton (*Gossypium hirsutum*), International Journal of Agriculture & Biology, 08–357/AKI/11–4–360–366.

Kawa A. Ali, Muhsin M. Qadir, Samira O. Rasool, Othman M. Hamad, (2012). The effect of spraying of wheat straw extracts on controlling some weed species, IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS), 1 : 36-39.

Kawa Abdulkareem Ali (2013). Allelopathic potential of some crop plant species on bread Wheat *Triticum aestivum* Using Equal compartment Agar Method, IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS), 2 : 52-55.

Khanh, T.D., Chung, M.I., Xuan, T.D. and Tawata, S., 2005. The exploitation of crop allelopathy in sustainable agricultural production. Journal of Agronomy and Crop Sciences 191, 172–184.

Lalitha, P1, Shivani K1, Rama Rao R1, (2012). *Parthenium hysterophorus* - an economical tool to increase the agricultural productivity, Int. J. Life Sci. Bt. & Pharm.

Leila Radhouane and Thouraya Rhim (2013). Allelopathic interaction of pepper (*Capsicum annuum*) and pearl millet (*Pennisetum glaucum*) intercropped, international journal of environment, 3 : 1.

M. Ayub1, M. Shehzad, M. A. Nadeem, M. Tahir, M. H. Siddiqui, M. Shoaib and I. Qadir (2013). Allelopathic effect of some summer legumes on seed germination, emergence and seedling growth of pearl millet (*Pennisetum americanum* L.), International Journal of Modern Agriculture, 2 : 4.

M.A. Randhawa, Z.A. Cheema and Muhammad Anjum Ali (2002). Allelopathic effect of sorghum water extract on the germination and seedling growth of *Trianthema portulacastrum*, International Journal of Agriculture & Biology, 1560–8530 /04–3–383–384.

M. B. Khan, M. Ahmad, M. Hussain, K. Jabran, S. Farooq, M. Waqas-Ul-Haq (2012). Allelopathic plant water extracts tank mixed with reduced doses of atrazine efficiently control *Trianthema portulacastrum* L. in *Zea mays L*., The Journal of Animal & Plant Sciences, 22(2) : 339-346.

Mahlagha Ghorbanli1, Afsaneh Gran and Amir Zolfaghary (2011). The study of Allelopathic potential in three species of *Glaucium* Mill. on *Sinapisarvensis* L. Iranian Journal of Plant Physiology, 2 (1), 321-325.

Mahmood A. Randhawa, Zahid A. Cheema & M. Saeed (1999). Mutual allelopathic effects of maize and *Trianthema portulacasturm* on germination and seedling growth, Pak. J. Agri. Sci. 36 (J -2).

Mahrokh Bolandi Amoghein, Roqiyeh Shakeri Amoghein, Ahmad Tobeh and Shahzad Jamaati-e-Somarin (2013). Allelopathic effects of extracts and plant residues of wild oat (*Avena fatua*) and rye (*Secale cereale* L.) on some germination parameters of wheat crop (*Triticum aestivum*) in the greenhouse condition, International Research Journal of Applied and Basic Sciences, 4 (8): 2313-2321.

Mohammed Worku (2010). Prevalence and distribution survey of an invasive alien weed (*Parthenium hysterophorus* L.) in Sheka zone, Southwestern Ethiopia, African Journal of Agricultural Research, 5(9): 922-927.

Molisch, H. (1937). Der Eingfluss einer Pflanze auf die andere- Allelopathie (Fischer, Jena).

Mudassir-ur-Rehman, Muhammad Ishaque, Ghulam Abbas, Muhammad Bakhsh Khokhar, Marghub Amer and Muhammad Irshad, (2013). Evaluation of combined application of allelopathic crop leachates and post-emergence herbicide for control of weeds in rice (*Oryza sativa* L.). Crop. Sci.Int. (Lahore), 25(4): 915-920.

Muhammad Ashraf and Muhammad Akhlaq (2007). Effects of sorghum leaves, roots and stems water extract, hand weeding and herbicide on weeds suppression and yield of wheat. Sarhad J. Agric., 23: 2.

Munesh Kumar and Sanjay Kumar (2010). Effect of *Parthenium hysterophorus* L. ash on growth and biomass of *Phaseolus mungo* L. Academia Arena, 2 : 2.

Muhammad Yasin, Zafar Iqbal, Muhammad Ehsan Safdar, Abdul Rehman, Amjed Ali, Muhammad Asif, Mudassir Aziz, Asif Tanveer and Muhammad Aslam Pervez (2011). *Phalaris minor* control, resistance development and strategies for integrated management of resistance to fenoxaprop-ethyl. African Journal of Biotechnology, 10(56) : 11802-11807.

Nasira Jabeen, Moinuddin Ahmed and Syed Shahid Shaukat (2011). Interactive activity of *asphodelus tenuifolius* on germination and growth of wheat (*Triticum aestivium* l.) and sorghum (*Sorghum bicolor* L.), Pak. J. Bot., 43(1) : 325-331.

P B S Bhadoria (2011). Allelopathy : A natural way towards weed management. American Journal of Experimental Agriculture, 1(1): 7-20.

P. K. Gantayet, K. C. Lenka and B. Padhy (2011). Vegetative growth and yield response of niger (*Guizotia abyssinica*) to leaf-litter dust of *Lantana camara*. Advances in Environmental Biology, 5(11): 3543-3549.

JR, Foy CL J Crop Prod 2001 ; 4:43 -119.

Rajan Katoch, Anita Singh, Neelam Thakur (2012). Allelopathic influence of dominant weeds of North-Western Himalayan region on common cereal crops. International Journal of Environmental Sciences, 3 : 1.

Rajan Katoch, Anita Singh and Neelam Thakur, (2012). Effect of weed residues on the physiology of common cereal crops. International Journal of Engineering Research and Applications (IJERA), 2(5): 828-834.

Rajiv P., Narendhran S., Subhash kumar M., Sankar A., Rajeshwari Sivaraj, Rajendran Venckatesh (2013). *Parthenium hysterophorus* L. compost: Assessment of its physical properties and allelopathic effect on germination and growth of *Arachis hypogeae* L. International Research Journal of Environment Sciences, 2(2) : 1-5.

Rahamdad Khan and Muhammad Azim Khan (2012). Weed control efficiency of bioherbicides and their impact on grain yield of wheat (*Triticum aestivum* L.). European Journal of Applied Sciences, 4 (5): 216-219.

Rice, E.L., 1984. Allelopathy. Academic Press, Orlando, 350 pp.

Rizvi, S.J.H., Haque, H., Singh, V.K. and Rizvi, V., 1992. A discipline called allelopathy, in Allelopathy: Basic and Applied Aspects, ed. by Rizvi SJH and Rizvi V. Chapman and Hall, London, UK, pp. 1–10.

Rezene Fessehaie, Mekasha Chichayibelu and Mengistu H. Giorgis, (2005). Spread and Ecological Consequences of *Parthenium hysterophorus* in Ethiopia. Arem., 6: 11-

Roth, C.M., Shroyer, J.P. and Paulsen, G.M., 2000. Allelopathy of sorghum on wheat under several tillage systems. Agronomy Journal 92, 885–860. *International Journal of Environment ISSN 2091-2854* 40 | P a g e

Saadia Bano, Muhammad Arshad Ullah, Abdul Khaliq, Khalida Hamid Abbasi and Saeeda Khanum (2012). Effect of aqueous extract of sun-dried neem (*Azadirachta indica*) leaves on wheat yield under field condition. Journal of Agricultural Extension and Rural Development, 4(12): 352-355.

S.B. Devaranavadgi, C. S. Hunshal, S.Y. Wali, M.B. Patil and M.A. Bellakki (2003). Studies on allelopathic effects in sorghum based agri-silviculture system. Karnataka J. Agril. Sci., 16(3) : 426-429.

Saheli Chatterjee, Sabyasachi Chatterjee, Aloke Bhattacharya and Sikha Dutta (2011). Allelopathic effect of *Cassia occidentalis* leaves on mustard seeds, An International Peer-Reviewed Journal, 1: 1.

Samad MA,Rahman MM, Hossain AKMM ,Rahman MS and RahmanSM J Soil Nature 2008 ; 2(2) : 13-18.

Santosh Kumar Mehar (2011). Assessment of effect of *Prosopis juliflora* litter extract on seed germination and growth of rice. Food Science and Quality Management, 2.

UK Sahoo (2013). Allelopathic studies of understorey weeds by agroforestry trees in home gardens of Mizoram. Journal of Experimental Biology and Agricultural Sciences, 1: 4.

Xuan, T.D., Shinkichi, T., Khanh, T.D. and Chung, I.M., 2005. Biological control of weeds and plant pathogens in paddy rice by exploiting plantallelopathy: An overview. Crop Protection 24(3), 197-206.

Z.A. Cheema, A. Khaliq (2000). Use of sorghum allelopathic properties to control weeds in irrigated wheat in a semi arid region of Punjab. Agriculture Ecosystems and Environment, 79 : 105–112.

Zimdahl , R. L.(1999). Fundamentals of Weed Science. 2nd ed. San Diego : Academic Press, pp. 450.